A NEW AUSTRALIAN SPECIES OF *DUBOISIA* R. BR. (SOLANACEAE)

L.A. Craven, B.J. Lepschi

Australian National Herbarium, CSIRO Division of Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia

&

L.A.R. Haegi

The Botanic Gardens of Adelaide and State Herbarium, North Terrace, Adelaide, South Australia 5000

Abstract

A new species of *Duboisia*, *D. arenitensis* Craven, Lepschi & Haegi, is described from the Northern Territory, Australia. The species is illustrated, its distribution mapped, and an identification key provided. The hyoscine and hyoscyamine content of the single collection known is 0.137% and 0.04%, respectively.

Introduction

The solanaceous genus *Duboisia* R. Br., consisting of three species, occurs in Australia and New Caledonia with one species, *D. myoporoides* R. Br., common to both areas. The other two species, *D. hopwoodii* (F. Muell.)F. Muell. and *D. leichhardtii* (F. Muell.)F. Muell., are restricted to Australia. The genus was treated comprehensively by Haegi (1983) and has been treated (for differing numbers of species) in several floras (e.g. Heine, 1976; Purdie *et al.*, 1982; Stanley and Ross, 1986; Conn, 1993). Species of the genus contain alkaloids and *D. myoporoides* and *D. leichhardtii–myoporoides* hybrids are cultivated for alkaloid production, notably for hyoscine and hyoscyamine. Plantations have been established in southeastern Queensland for this purpose. A concise account of the commercial utilisation of *Duboisia* in Australia has been given by Matheson (1979).

During a botanical exploration trip to western Arnhem Land, Northern Territory, in March 1984, a *Duboisia* was collected by L.A. Craven and G.M. Wightman on the floristically rich sandstone plateau in this region. The plant could not be identified with any of the known species of the genus and further study of the herbarium material has demonstrated that it possesses a unique combination of character states. It has been concluded that this material represents a new species of the genus, described here as *D. arenitensis*.

Duboisia arenitensis Craven, Lepschi & Haegi, sp. nov.

Frutex 1.5 m altus. Folia glabra; petiolo vel obsoleto usque 20 mm longo; lamina angustissime elliptica usque anguste ovata interdum ovati-elliptica, saepe falcata, 70×15 mm usque 100×25 mm. *Inflorescentia* paniculata, foliosa proximale, bracteata distale; pedicello 1.5–3.5 mm longo. Calyx 2.25–4.5 mm longus; lobis angustissime triangularibus usque sublinearibus, 1.25–3 mm longis, longitudine 0.6–0.7 calycis. Corolla 5–8 mm longa; lobis anguste ovatis usque anguste triangularibus, 1.75–4 mm longis, longitudine 0.4–0.5 corollae. Stamina 4, duobus 1–1.5 mm longis et ceteribus 2–2.5 mm longis, staminodio carenti. Fructus non visus.

Typus: Australia, Northern Territory, vicinity of Mt Gilruth, lat. 13°10'S, long. 133°06'E, 27.iii.1984, *Craven & Wightman 8292* (holo.: CANB; iso.: A, AD, DNA, E, G, K, L, MEL, MO, P, US).





Shrub 1.5 m tall, branches glabrous excepting some minute deciduous hairs at the nodes. *Leaves* glabrous; petiole obsolete or up to 20 mm long; lamina narrowly to very narrowly elliptic to narrowly ovate or occasionally ovate-elliptic, often falcate, 70×15 mm to 100×25 mm, progressively becoming reduced on flowering branchlets and then sublinear and straight in shape and 30×2 mm, the base attenuate, the apex long acuminate, the margin flat and entire to slightly and irregularly sinuate, the midrib prominent abaxially and adaxially plane or adaxially impressed proximally and prominulous distally. *Inflorescence* paniculate, leafy proximally, bracteate distally, with minute glandular hairs on the axes and bracts; pedicel glabrous or with sparse minute glandular hairs, 1.5-3.5 mm long, straight to slightly curved. *Calyx* with minute glandular hairs, these sparse on the abaxial surface and

more numerous on the adaxial surface, 2.25-4.5 mm long; lobes very narrowly triangular to sublinear, 1.25-3 mm long, 0.6-0.7 times as long as the whole calyx, the apex acute to subobtuse. *Corolla* with minute simple and minute glandular hairs on the abaxial surface and glabrous on the adaxial surface excepting the dense minute papillae on the distal portion of the lobes, 5-8 mm long; lobes narrowly ovate to narrowly triangular, 1.75-4 mm long, 0.4-0.5 times as long as the whole corolla, the apex acuminate to narrowly acute. *Stamens* 4, 2 of these 1–1.5 mm long and the others 2–2.5 mm long, staminode absent. *Gynoecium* 2–3 mm long. *Fruit* not seen. Fig. 1.

Distribution and ecology

The species is known only from the type gathering which was collected near Mt Gilruth on the western Arnhem Land plateau. Map 1.

The solitary plant from which this collection was made grew in shrubby woodland on sandy alluvium along a seasonal creek on the sandstone plateau. Five or six additional plants were observed growing nearby in a similar habitat.



Chemistry

Leaf material from the type collection has been analysed by Dr Bill Griffin, University of Queensland, Brisbane. The alkaloid content was low (hyo-scine 0.137%, hyo-scyamine 0.04%) (Griffin, pers. comm., 1990).

Although D. arenitensis may not warrant cultivating for its alkaloids alone, the species should be brought into cultivation for experimental purresearch poses. A program to obtain hybrids between the new and those species presently cultivated should be undertaken,

Map 1. Distributions of Duboisia species in Australia. D. arenitensis \triangle ; D. hopwoodii \bigcirc ; D. leichhardtii \circ ; D. myoporoides \lll .

and field research conducted to establish whether or not crop improvements could be made, either in alkaloid quality and/or quantity or in agronomic factors such as productivity, disease resistance, tolerance to harvesting, etc. Intergeneric and interspecific hybridisation is known to result in novel qualitative and/or quantitative chemical profiles in Solanaceae (Evans, 1986; El-Iman *et al.*, 1991) and the potential for *D. arenitensis* to contribute to crop improvement in *Duboisia* should be investigated.

L.A. Craven et al.

Notes

In the features reported in this paper, there is no single character state which distinguishes In the features reported in this paper, diete is no single character state which distinguishes D. arenitensis from its congeners; the species is separated on a combination of characters. D. arenuensis and the species are given in Table 1. The major differences between the species are given in Table 1.

Duboisia arenitensis not only extends the geographic range of the genus into the northern Duboisia dreinierists retritory (Map 1), but also extends its ecology. The species occurs part of the Northern zone of northern Australia where it is a component of the woodlands within the monsoon zone sandstone plateau of Arnham Lord, D.J. within the monstorr zensive sandstone plateau of Arnhem Land. Duboisia myoporoides developed on the extensive of eastern Australia usually in humid developed on the coastal region of eastern Australia, usually in humid eucalypt forests or occurs in the coastal region disturbed sites. Duboisia leichbardiii and a start of the in disturbed sites. occurs in the coastal ends often in disturbed sites. Duboisia leichhardtii occurs in open eucalypt forest rainforests and often in southcentral and southcentral of sout rainforests and order in southcentral and southeastern Queensland. The remaining and eucalypt woodlin is widely spread within the arid approximation of the remaining and eucalypt woodili, is widely spread within the arid zone and occurs in a range of species, *D. hopwoodili*, and multa woodlands communities including mallee and mulga woodlands.

The specific epithet is derived arbitrarily from the geological name for sandstone, arenite, and refers to the colloquial name used by botanists for sandstone plateau country of northern Australia, i.e. "the sandstone".

	D. arenitensis	D. hopwoodii	D. leichhardtii	D. myoporoides
Leaves	Usually petiolate, rarely subsessile or sessile. Petiole 1–20 mm long.	Sessile to subsessile, rarely petiolate. Petiole 3–10 mm long.	Sessile to petiolate. Petiole 1–8 mm long.	Sessile to petiolate. Petiole 1–30 mm long.
Lamina	Narrowly elliptic to narrowly ovate, rarely sublinear (on flowering branchlets). Often falcate.	Narrowly to very narrowly elliptic to linear.	Narrowly to very narrowly ovate-elliptic, rarely narrowly ovate or narrowly elliptic.	Narrowly obovate to narrowly obovate- elliptic.
Lomina length	30-100 mm	24–125 mm	51–135 mm	38–150 mm
Lamina tengui Lamina breadth Lamina apex	2-25 mm	1–13 mm	7–22 mm	1–43 mm
	Long acuminate	Narrowly acute to acuminate, rarely apiculate.	Acuminate or narrowly acute.	Blunt, acute to obtuse, rarely rounded.
Pedicel length (in flower)	1.5–3.5 mm	1.5–5 mm	10–16 mm	2–7 mm
Calvx length	2.25-4.5 mm	1.5-4.5 mm	1.5-3.2 mm	1.2–3.2 mm
Calyx lobe shape	Very narrowly triangular to sublinear	Triangular to broadly triangular, occasionally acuminate.	Very narrowly triangular to more or less linear.	Triangular to broadly triangular.
Calvx lobe length	1.25–3 mm	0,3–2 mm	0.5–1.5 mm	0.2-0.8 mm
Calyx lobe length: overall calyx length.	0.6-0.7	0.3-0.8	0.5-0.6	0.1-0.3
Corolla length	5–8 mm	7–15.5 mm	13–19 mm	47 mm
Corolla lobe shape	Narrowly ovate to narrowly triangular.	Broadly ovate to ovate-elliptic or orbicular.	Very narrowly triangular to linear.	Broadly ovate to ovate-elliptic or rarely orbicular.
Corolla lobe length:	1.75–4 mm	1–5 mm	6.5-11.5 mm	1.3→3.5 mm
Corolla lobe length, overall corolla length	0.4–0.5	0.1-0.3	0.5-0.6	0.3-0.5

Table 1 - Major distinguishing characters of Duboisia species

J. Adelaide Bot. Gard. 16: 27-31 (1995)

Duboisia arenitensis (Solanaceae)

Key to the species of *Duboisia*

1. Leaves with the lamina mostly narrowly elliptic to linear.

- Leaves petiolate, rarely subsessile or sessile. Calyx lobes very narrowly triangular to sublinear. Corolla lobes narrowly ovate to narrowly triangular, 0.4–0.5 times as long as the corolla; corolla 5–8 mm long D. arenitensis
- Leaves sessile or subsessile, rarely petiolate. Calyx lobes triangular to broadly triangular. Corolla lobes broadly ovate to orbicular, 0.1-0.3 times as long as the corolla; corolla 7-15.5 mm long

D. hopwoodii 1. Leaves with the lamina mostly obovate to obovate-elliptic or ovate to narrowly ovate or ovate-elliptic.

- Leaves with the lamina mostly ovate to narrowly ovate or ovate-elliptic, apex acuminate. Corolla lobes narrowly acute.
 - 4. Pedicel 1.5-3.5 mm long, Corolla 5-8 mm long, corolla lobes 1.75-4 mm longD. arenitensis
 - 4. Pedicel 10-16 mm long. Corolla 13-19 mm long, corolla lobes 6.5-11.5 mm longD. leichhardtii

Acknowledgements

Bill Griffin is thanked for carrying out the alkaloid analysis of *D. arenitensis*. Catherine Wardrop prepared the illustration.

References

Conn, B. (1993). Solanaceae. In Harden, G.J. (ed.). "Flora of New South Wales", Vol. 3. (University of New South Wales Press: Sydney).

El-Imam, Y.M.A., Evans, W.C., Haegi, L., & Ramsey, K.P.A. (1991). Secondary metabolites of intergeneric hybrids of the Anthocercideae, family Solanaceae. Int. J. Pharmacognosy, 29, 263–267.

Evans, W.C. (1986). Hybridisation and secondary metabolism in the Solanaceae. In D'Arcy, W.G. (ed.). "Solanaceae: Biology and Systematics". (Columbia University Press: New York).

Haegi, L.A.R. (1983). "Systematic and Evolutionary Studies in the Australian Solanaceae". Ph.D. thesis, University of Adelaide.

 Heine, H. (1976). Solanaceae. In Aubréville, A. and Leroy, J-F. (eds). "Flore de la Nouvelle-Calédonie et Dépendances", Vol. 7. (Muséum National d'Histoire Naturelle: Paris).
Matheson, E.M. (1979). Other crops of potential importance. In Lovett, J.V. and Lazenby, A. (eds). "Australian

Matheson, E.M. (1979). Other crops of potential importance. In Lovett, J.V. and Lazenby, A. (eds). "Australian Field Crops. Vol. 2. Tropical Cerals, Oilseeds, Grain Legumes and Other Crops". (Angus & Robertson: Sydney).

Purdie, R.W., Symon, D.E. & Haegi, L.A.R. (1982). Solanaceae. In George, A.S. (ed.). "Flora of Australia", Vol. 29. (Australian Government Publishing Service: Canberra).

Stanley, T.D. & Ross, E.M. (1986). Solanaceae. "Flora of South Eastern Queensland", Vol. 2. (Queensland Department of Primary Industries: Brisbane).